

Supporting Information for

Color-tunable hetero-dinuclear Pt(II)/B(III) and
Pt(II)/Ir(III) arrays with N[^]O-julolidine ligands

Adela Nano,^{,a} Maria Pia Gullo,^b Barbara Ventura,^b Andrea Barbieri,^{*,b} Nicola Armaroli,^b
Raymond Ziessel^a*

^a Institut de Chimie et Procédés pour l'Énergie, l'Environnement et la Santé (ICPEES), UMR
7515 au CNRS, Ecole Européenne de Chimie, Polymères et Matériaux (ECPM), 25 rue
Becquerel, 67087 Strasbourg Cedex 02, FRANCE.

^b Istituto per la Sintesi Organica e la Fotoreattività (ISO-F), Consiglio Nazionale delle Ricerche
(CNR), Via P. Gobetti 101, 40129 Bologna BO, ITALY.

Table of Contents

| | |
|---|-----|
| Figure S1. ^1H -NMR spectrum of PtL₃ in CDCl_3 | S3 |
| Figure S2. ^{13}C -NMR spectrum of PtL₃ in CDCl_3 | S4 |
| Figure S3. ^1H -NMR spectrum of PtBL₃ in CDCl_3 | S5 |
| Figure S4. ^{13}C -NMR spectrum of PtBL₃ in CDCl_3 | S6 |
| Figure S5. ^1H -NMR spectrum of IrL₁ in CDCl_3 | S7 |
| Figure S6. ^{13}C -NMR spectrum of IrL₁ in CDCl_3 | S8 |
| Figure S7. ^1H -NMR spectrum of IrL₂ in C_6D_6 | S9 |
| Figure S8. ^{13}C -NMR spectrum of IrL₂ in C_6D_6 | S10 |
| Figure S9. ^1H -NMR spectrum of IrL₃ in C_6D_6 | S11 |
| Figure S10. ^{13}C -NMR spectrum of IrL₃ in C_6D_6 | S12 |
| Figure S11. ^1H -NMR spectrum of PtIrL₃ in CD_3CN | S13 |
| Figure S12. ^{13}C -NMR spectrum of PtIrL₃ in CD_3CN | S14 |
| Figure S13. Absorption spectra of IrL₁ and IrL₃ and the relevant models at rt | S15 |
| Figure S14. Absorption spectra of PtL₃ and PtBL₃ and the relevant models at rt | S16 |
| Figure S15. Absorption spectra of PtIrL₃ and of the relevant models at rt | S17 |
| Figure S16. Emission/Excitation contour map of PtIrL₃ at 77 K | S18 |




Figure S1. 1H -NMR spectrum of PtL_3 in $CDCl_3$




Figure S2. ^{13}C -NMR spectrum of PtL_3 in CDCl_3




Figure S3. $^1\text{H-NMR}$ spectrum of PtBL_3 in CDCl_3




Figure S4. ¹³C-NMR spectrum of **PtBL₃** in CDCl₃




Figure S6. ¹³C-NMR spectrum of IrL₁ in CDCl₃




Figure S7. ¹H-NMR spectrum of IrL₂ in C₆D₆





Figure S8. ¹³C-NMR spectrum of **IrL₂** in C₆D₆






Figure S10. ¹³C-NMR spectrum of **IrL₃** in C₆D₆




Figure S11. ^1H -NMR spectrum of PtIrL_3 in CD_3CN




Figure S12. ^{13}C -NMR spectrum of PtIrL_3 in CD_3CN




Figure S13. Absorption spectra of IrL_1 and the relevant models Ir and L_3 (top), and of IrL_3 and the relevant models Ir and L_3 (bottom) in CH_3CN solution at rt. The dotted line represents the spectral addition of the model components.




Figure S14. Absorption spectra of the array PtL_3 and the relevant models Pt and L_3 (top), and of the array $PtBL_3$ and the relevant models Pt and BL_3 (bottom) in CH_3CN solution at rt. The dotted line represents the spectral addition of the model components.




Figure S15. Absorption spectra of the multichromophoric system **PtIrL₃** and the relevant models **Pt** and **IrL₃** in CH_3CN solution at rt. The dotted line represents the spectral addition of the model components.




Figure S16. Emission/Excitation contour map of the multichromophoric system **PtIrL₃** in $\text{CH}_3\text{OH}:\text{C}_2\text{H}_5\text{OH}$ (1:4) mixture at 77 K.